



Slopes of Siegel cusp forms and geometry of compactified Kuga varieties

Flora Poon, Riccardo Salvati Manni and Gregory Sankaran

To Gerard, with friendship and admiration

ABSTRACT

We study the Kodaira dimension of the compactified n -fold Kuga variety over the moduli space of principally polarised abelian g -folds. We construct a suitable compactification, which we call a Namikawa compactification, and show that in most cases it has canonical singularities. We then use results about the slope of Siegel modular forms to determine the Kodaira dimension for all $g > 1$ and $n > 0$.

1. Introduction

The universal family $X(\Gamma)$ over a moduli space $\mathcal{A}(\Gamma) = \Gamma \backslash \mathbb{H}_g$ of abelian varieties associated with a finite-index subgroup Γ of $\mathrm{Sp}(2g, \mathbb{Z})$ is known as the Kuga variety. Such families were first studied systematically by Kuga, whose 1964 Chicago lecture notes on the subject [Kug18] have been recently published. The construction is given in [Kug18] for $\Gamma < \mathrm{Sp}(2g, \mathbb{Z})$ a torsion-free subgroup of finite index, but the restriction to torsion-free can be removed. However, if $-1 \in \Gamma$, then the fibre of the family is no longer the abelian variety but instead the corresponding Kummer variety. One could also allow Γ to be a subgroup of $\mathrm{Sp}(2g, \mathbb{Q})$ commensurable with $\mathrm{Sp}(2g, \mathbb{Z})$, for example taking $\mathcal{A}(\Gamma)$ to be the moduli space of abelian varieties with some non-principal polarisation, but we shall not pursue this here.

A natural generalisation is to consider the n -fold Kuga variety $X^n(\Gamma)$, whose general fibre is the n -fold direct product A^n of the corresponding abelian variety A or, if $-1 \in \Gamma$, the Kummer variety $A^n / \pm 1$. Alternatively, one may consider the universal family $\underline{X}^n(\Gamma)$ over the stack $\underline{\mathcal{A}}(\Gamma) := [\Gamma \backslash \mathbb{H}_g]$. In this case the fibre is an abelian variety in all cases, but if $-1 \in \Gamma$, then the base has non-trivial stabilisers generically. This is the object that is studied in a particular case in [FV16].

We shall be concerned with compactifications of the n -fold Kuga variety $X_g^n = X^n(\mathrm{Sp}(2g, \mathbb{Z}))$ associated with the coarse moduli space \mathcal{A}_g of principally polarised abelian g -folds over \mathbb{C} . Thus $X_g^1 = X(\mathrm{Sp}(2g, \mathbb{Z}))$, and by convention $X_g^0 = \mathcal{A}_g$.

The starting point for our work is Ma's study [Ma21] of $X_g^n(\Gamma)$. We construct a special compactification, which we call a Namikawa compactification, of X_g^n , and this, together with recent

Received 14 September 2021, accepted in final form 5 July 2025.

2020 Mathematics Subject Classification 14K10 (primary), 11F46 (secondary).

Keywords: Kodaira dimension, Kuga variety, Siegel modular forms, Toroidal compactification.

This journal is © Foundation Compositio Mathematica 2025. This article is distributed with Open Access under the terms of the [Creative Commons Attribution Non-Commercial License](#), which permits non-commercial reuse, distribution, and reproduction in any medium, provided that the original work is properly cited. For commercial re-use, please contact the [Foundation Compositio Mathematica](#).

and less recent results about the slope of \mathcal{A}_g , allows us to determine the Kodaira dimension $\kappa(X_g^n)$ of X_g^n whenever $g \geq 2$ and $n \geq 1$.

THEOREM 1. *Suppose $g \geq 2$ and $n \geq 1$. Then the Kodaira dimension $\kappa(X_g^n)$ of X_g^n satisfies*

- $\kappa(X_g^n) = \frac{1}{2}g(g+1) = \dim \mathcal{A}_g$ if $g+n \geq 7$, except for $(g, n) \in \{(4, 3), (3, 5), (3, 4), (2, 7), (2, 6), (2, 5)\}$;
- $\kappa(X_2^7) = \kappa(X_3^5) = \kappa(X_4^3) = 0$; and
- $\kappa(X_g^n) = -\infty$ otherwise, that is, if $g+n \leq 6$ or $(g, n) \in \{(2, 6), (2, 5), (3, 4)\}$.

For completeness, we mention the case $g = 1$. We can identify X_1^n with $\mathcal{M}_{1, n+1}$, the coarse moduli space of genus 1 curves with $n+1$ (ordered) marked points: the case $n = 0$ is the familiar identification of $\mathcal{M}_{1,1}$ with \mathbb{A}^1 via the j -invariant. The results of Belorousski [Bel98] then show that X_1^n is rational when $n \leq 9$, and from [BF06] we have $\kappa(X_1^{10}) = 0$ and $\kappa(X_1^n) = 1$ for $n \geq 11$.

In fact, X_2^n for $n \leq 6$ and X_3^n for $n \leq 4$ are all unirational. If $g \leq 3$, then, since \mathcal{A}_g is birationally equivalent to \mathcal{M}_g and the g th symmetric power of a curve is birational to its Jacobian, there is a dominating rational map $\mathcal{M}_{g, gn} \dashrightarrow X_g^n$. On the other hand, it is known that $\mathcal{M}_{2,n}$ and $\mathcal{M}_{3,n}$ are unirational for $n < 13$ and $n < 15$, respectively; see [KT21, Table 1].

Moreover, X_g^1 is unirational for $g \leq 5$; see [Ver08] for $g = 3$ and $g = 4$ and [FV16] for $g = 5$. Note that X_g^1 is the boundary of the Mumford partial compactification \mathcal{A}'_{g+1} . One can therefore compactify X_g^1 by taking its closure in any toroidal compactification of \mathcal{A}_{g+1} since these all contain \mathcal{A}'_{g+1} as a dense open set, but it is not straightforward to control the singularities that then arise.

For the cases $g+n \leq 6$, our proof is based on the fact that the slope of \mathcal{A}_g is greater than 7: this was proved in [Sal92] for $g = 3$ and $g = 4$ and in [FGSV14] for $g = 5$. For the cases $g+n \geq 7$, our main technical result concerns the existence of a sufficiently good compactification of X_g^n . We will say that a compactification of X_g^n is a Namikawa compactification if it dominates a toroidal compactification of \mathcal{A}_g and boundary divisors are mapped to boundary divisors; see Definition 2.1 for a full explanation and precise details. We prove the following (see Theorem 2.2 for the precise statement).

THEOREM 2. *Suppose $g \geq 2$ and $n \geq 1$. There exists a Namikawa compactification $\overline{X_g^n}$ with canonical singularities as long as $g+n \geq 6$.*

There is some overlap between our results and those of [Ma21]. The singularities of X_g^n are studied in [Ma21, § 10], but the singularities at the boundary $\overline{X_g^n}$ are not considered there. That is sufficient for computing the geometric genus but not other plurigenera; on the other hand, Ma's approach gives precise information about the geometric genus and therefore some information about the Kodaira dimension.

Our original motivation came from the case $g = 6$ and $n = 1$ and the Kodaira dimension $\kappa(X_6^1)$. As we have seen, X_g^1 is unirational for $g < 6$. On the other hand, for $g \geq 7$ we have $\kappa(X_g^1) = \kappa(\mathcal{A}_g) = \frac{1}{2}g(g+1)$, since Iitaka's conjecture holds by [Kaw85] and \mathcal{A}_g is of general type.

Similar questions arise in relation to the universal Jacobian varieties over \mathcal{M}_g . A suitable compactification was constructed by Caporaso [Cap94], and the question of Kodaira dimension is well studied; see for example [BFV12, CKV17, FV13], as well as [BF06] for the case $g = 1$.

2. Namikawa compactifications

Suppose throughout this section that $g \geq 2$. Let $\mathcal{A}_g \subset \overline{\mathcal{A}}_g$ be the inclusion of the coarse moduli space $\mathcal{A}_g = \mathrm{Sp}(2g, \mathbb{Z}) \backslash \mathbb{H}_g$ of principally polarised abelian varieties of dimension g into a toroidal compactification $\overline{\mathcal{A}}_g$. Denote by $f: X_g^n = (\mathbb{Z}^{2gn} \ltimes \mathrm{Sp}(2g, \mathbb{Z})) \backslash (\mathbb{C}^{gn} \times \mathbb{H}_g) \rightarrow \mathcal{A}_g$ the n -fold Kuga family, as in [Ma21].

DEFINITION 2.1. A *Namikawa compactification* of X_g^n is an irreducible normal projective variety $\overline{X_g^n}$ containing X_g^n as an open subset, together with a projective toroidal compactification $\overline{\mathcal{A}}_g$ of \mathcal{A}_g for which the following conditions hold:

- (i) The morphism $f: X_g^n \rightarrow \mathcal{A}_g$ extends to a projective morphism $f: \overline{X_g^n} \rightarrow \overline{\mathcal{A}}_g$.
- (ii) Every irreducible component of $\Delta_X := \overline{X_g^n} \setminus X_g^n$ dominates an irreducible component of $\Delta_{\mathcal{A}} := \overline{\mathcal{A}}_g \setminus \mathcal{A}_g$.

Compactifications satisfying these conditions were first given by Namikawa [Nam79, Nam80], for the case $n = 1$. There is little difficulty in extending Namikawa's construction to arbitrary n , and it is essentially done in [FC90, § VI.1]. The conditions in Definition 2.1 are the same as in [Ma21, Theorem 1.2], but we also require $\overline{X_g^n}$ to be normal (rather than just smooth in codimension 1) and projective. In fact, neither of these conditions presents any difficulty.

THEOREM 2.2. Suppose $g \geq 2$ and $n \geq 1$. Then if $g + n \geq 6$, there exists a Namikawa compactification $\overline{X_g^n} \supset X_g^n$ such that $\overline{X_g^n}$ has canonical singularities. In particular, $\overline{X_g^n}$ is \mathbb{Q} -Gorenstein.

The proof will occupy the rest of this section. The first step is the following lemma. Although similar statements are already known (see for example [AS16]), we give a proof here as we have not found one elsewhere.

Suppose that a finite group G acts effectively on a variety X . A non-trivial element of G is called a *quasireflection* if it preserves a divisor on X .

LEMMA 2.3. Suppose that G is a finite group acting effectively and without quasireflections on a variety X that has canonical singularities. Let $f: \hat{X} \rightarrow X$ be a G -equivariant resolution of singularities, and suppose that \hat{X}/G has canonical singularities. Then X/G also has canonical singularities.

Proof. Since X has canonical singularities, it is in particular \mathbb{Q} -Gorenstein (we do not require, nor expect, X or X/G to be \mathbb{Q} -factorial), and therefore X/G is also \mathbb{Q} -Gorenstein. Suppose that $rK_{X/G}$ is Cartier and $\sigma \in \mathcal{O}(rK_{X/G})$, so σ is a pluricanonical form on $(X/G)_{\mathrm{reg}}$. Therefore, σ lifts to a G -invariant form $g^*\sigma$ on an open G -invariant subset $X_0 \subset X$, where $g: X \rightarrow X/G$ is the quotient map.

The complement $X \setminus X_0$ consists of the fixed loci of elements of G , together with the singular locus of X ; however, the fixed loci have codimension at least 2 by assumption, so $g^*\sigma$ extends G -invariantly to X_{reg} . Therefore, it lifts on \hat{X} to a form defined away from the exceptional locus of f . As X has canonical singularities, this extends to a G -invariant form $\hat{\sigma} = f^*g^*\sigma$ on \hat{X} without poles, which in turn descends to the smooth part $(\hat{X}/G)_{\mathrm{reg}}$ of \hat{X}/G because $\hat{g}: \hat{X} \rightarrow \hat{X}/G$ is étale in codimension 1.

This form $\hat{\sigma}$ agrees with $\bar{f}^*\sigma$ on the dense open set $\bar{f}^{-1}((X/G)_{\mathrm{reg}})$, where $\bar{f}: \hat{X}/G \rightarrow X/G$ is the map induced by f ; therefore, $\bar{f}^*\sigma$ extends to $(\hat{X}/G)_{\mathrm{reg}}$.

Now consider a resolution of singularities $h: Y \rightarrow X/G$, and form the pullback $\hat{h}: \hat{Y} \rightarrow \hat{X}/G$ as in the diagram

$$\begin{array}{ccccc}
 \hat{X} & \xrightarrow{f} & X & & \\
 \hat{g} \searrow & & & \swarrow g & \\
 & \hat{X}/G & \xrightarrow{\bar{f}} & X/G & \\
 \hat{h} \nearrow & & & \nearrow h & \\
 \hat{Y} & \xrightarrow{\hat{f}} & Y & &
 \end{array}$$

This resolves the singularities of \hat{X}/G , so $\hat{h}^* \bar{f}^* \sigma$ extends without poles to the whole of \hat{Y} . But now if $h^* \sigma$ has poles along a divisor $E \subset Y$, then $\hat{f}^* h^* \sigma = \hat{h}^* \bar{f}^* \sigma$ has poles along $\hat{f}^{-1} E$, which is impossible. Therefore, $h^* \sigma$ is holomorphic, and hence X/G has canonical singularities. \square

Next, we construct Namikawa compactifications $\overline{X_g^n}$. To do this we follow [Nam79] closely, and, as far as possible, we have chosen our notation to be compatible with that paper. The notation in [FC90] is different. Note that our compactifications are singular; both [Nam79, Proposition 5.4(i)] and [FC90, Theorem VI.1.1(i)] mention smooth compactifications, but after a base change in the first case and as stacks in the second (see [FC90, § VI.1, top of p. 195]).

We write $\Gamma = \mathrm{Sp}(2g, \mathbb{Z})$. For simplicity, and because it is enough for our purposes, we consider only this case, but allowing Γ to be a finite-index subgroup of $\mathrm{Sp}(2g, \mathbb{Z})$ does not change the argument. To fix notation, we choose a free \mathbb{Z} -module \mathbb{W} of rank $2g$ equipped with a \mathbb{Z} -basis e_1, \dots, e_{2g} , and fix a standard skew-symmetric form by requiring $\langle e_i, e_{i+g} \rangle = -1$ for $1 \leq i \leq g$ and $\langle e_i, e_j \rangle = 0$ if $|i - j| \neq g$, so that the matrix of the form with respect to the basis $\{e_i\}$ is $\begin{pmatrix} 0 & -\mathbf{1}_g \\ \mathbf{1}_g & 0 \end{pmatrix}$ (we denote the $r \times r$ identity matrix by $\mathbf{1}_r$ throughout). Then

$$\mathrm{Sp}(2g, \mathbb{Z}) = \left\{ \gamma \in \mathrm{GL}(2g, \mathbb{Z}) \mid {}^t \gamma \begin{pmatrix} 0 & -\mathbf{1}_g \\ \mathbf{1}_g & 0 \end{pmatrix} \gamma = \begin{pmatrix} 0 & -\mathbf{1}_g \\ \mathbf{1}_g & 0 \end{pmatrix} \right\}.$$

Choose a cusp $F_{g'}$ of rank g' of \mathbb{H}_g for some $0 \leq g' < g$, and put $g'' = g - g'$. Recall that \mathbb{H}_g is the cone of symmetric $g \times g$ complex matrices M with positive-definite imaginary part: cusps of rank g' arise by allowing $\mathrm{Im} M$ to be semi-definite but requiring the radical $\mathrm{rad} M$ to be defined over \mathbb{Q} and of dimension g'' . Such a cusp thus corresponds to a rank g'' isotropic sublattice \mathbb{X} of \mathbb{W} up to the action of Γ , but $\Gamma = \mathrm{Sp}(2g, \mathbb{Z})$ acts transitively on such lattices. Therefore, without loss of generality, we may take $F_{g'}$ to have stabiliser

$$P(g') = \left\{ \begin{pmatrix} A & 0 & B & m' \\ m & u & n & M \\ C & 0 & D & n' \\ 0 & 0 & 0 & {}^t u^{-1} \end{pmatrix} \mid \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}(2g', \mathbb{R}), u \in \mathrm{GL}(g'', \mathbb{R}) \right\}$$

in $\mathrm{Sp}(2g, \mathbb{R})$, by choosing $\mathbb{X} = \mathbb{Z}e_{g'+1} + \dots + \mathbb{Z}e_g \cong \mathbb{Z}^{g''}$.

Next we consider the integral affine symplectic group $\tilde{\Gamma}_g^n$. It is given by

$$\tilde{\Gamma}^n = \mathbb{Z}^{2gn} \ltimes \mathrm{Sp}(2g, \mathbb{Z}) < \tilde{\Gamma}_{\mathbb{R}}^n = \mathbb{R}^{2gn} \ltimes \mathrm{Sp}(2g, \mathbb{R}) < \mathrm{GL}(n+2g, \mathbb{R})$$

and consists of elements $\tilde{\gamma}$ of the form

$$\tilde{\gamma} = \begin{pmatrix} 1 & a & b \\ 0 & A_0 & B_0 \\ 0 & C_0 & D_0 \end{pmatrix}, \quad \gamma = \begin{pmatrix} A_0 & B_0 \\ C_0 & D_0 \end{pmatrix} \in \mathrm{Sp}(2g, \mathbb{R}), \quad a, b \in M_{n \times g}(\mathbb{R}) \quad (2.1)$$

(cf. [Nam79, § 2.7]).

The integral affine symplectic group acts on $\mathbb{C}^{gn} \times \mathbb{H}_g$, and the Kuga variety, cf. [Nam79, equation (3.4.1)], is the quotient

$$X_g^n := \tilde{\Gamma}^n \backslash (\mathbb{C}^{gn} \times \mathbb{H}_g).$$

It is the coarse moduli space of principally polarised abelian varieties together with a point on the Kummer variety of the n -fold cartesian self-product.

The stabiliser of $F_{g'}$ in $\tilde{\Gamma}_{\mathbb{R}}^n$ is (cf. [Nam79, Example (2.8)])

$$\tilde{P}(g') = \left\{ \begin{pmatrix} \mathbf{1}_n & a' & a'' & b' & b'' \\ 0 & A & 0 & B & m' \\ 0 & m & u & n & M \\ 0 & C & 0 & D & n' \\ 0 & 0 & 0 & 0 & {}^t u^{-1} \end{pmatrix} \middle| \begin{array}{l} \gamma' = \begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \mathrm{Sp}(2g', \mathbb{R}), \\ u \in \mathrm{GL}(g'', \mathbb{R}), \\ a', b' \in M_{n \times g'}(\mathbb{R}), \\ a'', b'' \in M_{n \times g''}(\mathbb{R}) \end{array} \right\}, \quad (2.2)$$

where as before M , m and n , and m' and n' are subject to the symplecticity conditions, and its unipotent radical has centre

$$\tilde{U}(g') = \{u(b'', M) \mid M = {}^t M\}, \quad \text{where } u(b'', M) = \begin{pmatrix} 1 & 0 & 0 & 0 & b'' \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & M \\ 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 1 \end{pmatrix}.$$

Intersecting with $\tilde{\Gamma}^n$, we obtain the group

$$\tilde{\Upsilon}^n = \tilde{U}(g') \cap \tilde{P}(g') = \{u(b'', M) \mid b'' \in \mathbb{Z}^{ng''}, M = {}^t M \in M_{g'' \times g''}(\mathbb{Z})\},$$

which is identified with $\mathrm{Sym}^2(\mathbb{X}^\vee) \times (\mathbb{X}^\vee)^n$. (This means a symmetric bilinear function on \mathbb{X} and n linear functions on \mathbb{X} . In other words, $\mathrm{Sym}^1(\mathbb{X}^\vee)$ is \mathbb{X}^\vee . It is what is called $B(X) \oplus (X^*)^n$ in [FC90].)

To obtain the partial compactification at the cusp $F_{g'}$, we first take the partial quotient by $\tilde{\Upsilon}^n$. For this we use the Siegel domain realisation of \mathbb{H}_g : for $\tau \in \mathbb{H}_g$, we write

$$\tau = \begin{pmatrix} \tau' & \omega \\ {}^t \omega & \tau'' \end{pmatrix}$$

with $\tau' \in \mathbb{H}_{g'}$, $\omega \in M_{g' \times g''}(\mathbb{C})$ and $\tau'' \in M_{g'' \times g''}^{\mathrm{sym}}(\mathbb{C})$, and then

$$\mathbb{H}_g \cong \mathcal{D}_{g'} := \{(\tau', \omega, \tau'') \mid \mathrm{Im} \tau'' - (\mathrm{Im} {}^t \omega)(\mathrm{Im} \tau')^{-1}(\mathrm{Im} \omega) > 0\}.$$

Then $M \in \mathrm{Sym}^2(\mathbb{X})$ acts by translations in the imaginary directions in $M_{g'' \times g''}^{\mathrm{sym}}(\mathbb{C})$, so near this boundary, X_g^n is covered by

$$\mathcal{D}_{g'} \times \mathbb{C}^{ng'} \times (\mathbb{C}^*)^{ng''} \subset \mathbb{H}_{g'} \times \mathbb{C}^{g'g''} \times (\mathbb{C}^*)_{\mathrm{sym}}^{g'' \times g''} \times (\mathbb{C}^{g'} \times (\mathbb{C}^*)^{g''})^n, \quad (2.3)$$

where the term $((\mathbb{C}^*)^{g''})^n$ is $\mathbb{C}^{ng''}/(\mathbb{X}^\vee)^n$, given by b'' acting by translations in the imaginary directions in $\mathbb{C}^{ng''} = \mathbb{X}^\vee \otimes \mathbb{C}$.

Now we compactify by replacing the torus part $(\mathbb{C}^*)_{\text{sym}}^{g'' \times g''} \times ((\mathbb{C}^*)^{g''})^n$ (that is, the quotient $\mathbb{C}^{(g''^2 + (2n+1)g'')/2} / \tilde{\Upsilon}^n$) with a suitable torus embedding $\text{Temb}(\Sigma(g'))$ corresponding to a fan $\Sigma(g')$ in $(\text{Sym}^2(\mathbb{X}^\vee) \times (\mathbb{X}^\vee)^n) \otimes \mathbb{R}$. We first define two (non-polyhedral) cones, in $\text{Sym}^2(\mathbb{X}^\vee)$ and $\text{Sym}^2(\mathbb{X}^\vee) \times (\mathbb{X}^\vee)^n$, respectively, following [FC90].

The first cone is $C(\mathbb{X}) \subset \text{Sym}^2(\mathbb{X}^\vee \otimes \mathbb{R})$, which is defined to be the cone of positive semi-definite symmetric bilinear forms b on $\mathbb{X}^\vee \otimes \mathbb{R}$ with rational radical $\text{rad } b$. Equivalently, $C(\mathbb{X})$ is the cone generated by the rank 1 forms in $\text{Sym}^2(\mathbb{X}^\vee \otimes \mathbb{Q})$. To construct a toroidal compactification of \mathcal{A}_g , one must give decompositions of these cones for each $g_i < g$ into fans invariant under the stabiliser in Γ of the cusp $F(g')$. There are many ways to do that, but using reduction theory of quadratic forms (which guarantees the compatibility) supplies two important ones, the first and second Voronoi decomposition.

The second cone is $\tilde{C}(\mathbb{X}) \subset (\text{Sym}^2(\mathbb{X}^\vee) \oplus (\mathbb{X}^\vee)^n) \otimes \mathbb{R}$, given by

$$\tilde{C}(\mathbb{X}) = \{(b, \ell_1, \dots, \ell_n) \in \text{Sym}^2(\mathbb{X}^\vee \otimes \mathbb{R}) \oplus (\mathbb{X}^\vee \otimes \mathbb{R})^n \mid \ell_i|_{\text{rad } b} = 0 \text{ for all } i\},$$

as described in [FC90, Definition VI.1.3]. The number n does not change, so we suppress it in the notation, but observe that $\tilde{C}(\mathbb{X})$ implicitly depends on n whereas $C(\mathbb{X})$ does not.

We then further decompose $\tilde{C}(\mathbb{X})$ as in [FC90, Definition VI.1.3], for each g' , obtaining a collection of fans $\tilde{\Sigma} = \{\tilde{\Sigma}(g') \mid 0 \leq g' < g\}$. Provided that we choose the fans compatibly for different cusps, we obtain, by the standard toroidal compactification procedure from [AMRT10], a compactification $\overline{X_g^n}^{\tilde{\Sigma}}$ that is analytically locally isomorphic to the product of a smooth space with a quotient of $\text{Temb}(\tilde{\Sigma}(g'))$ by $\tilde{\Gamma}^n \cap P(g')$. Ultimately, we shall choose $\tilde{\Sigma}(g')$ to be regular so that $\text{Temb}(\tilde{\Sigma}(g'))$ is smooth. Singularities will then arise when we move beyond $\tilde{\Upsilon}^n$ and take the quotient by the rest of $\tilde{\Gamma}^n$, which in our situation may have fixed points.

We need more, however, because we seek a Namikawa compactification. These are alluded to in [FC90], but the equidimensional condition [FC90, Definition VI.1.3(v)], which is not used and therefore not examined in detail there, is crucial for us. To be precise, we require a weaker version of equidimensional, which we call equidimensional in codimension 1. For this, it is enough if every ray in $\tilde{\Sigma}(g')$ maps onto a cone of the fans Σ that define $\overline{\mathcal{A}}_g$, whereas equidimensionality requires this for every cone in $\tilde{\Sigma}(g')$ of any dimension.

These two conditions, smoothness and equidimensionality, are in general opposed to one another. Choosing $\tilde{\Sigma}$ to give smooth covering spaces $\text{Temb}(\tilde{\Sigma}(g'))$ typically involves blowing up, and thus instantly violates condition (ii) of Definition 2.1. Therefore, to construct an appropriate $\overline{X_g^n}$, we need a slightly more indirect approach. Instead of taking a regular decomposition straight away, we first choose a decomposition $\tilde{\Sigma}^\flat$ such that $\text{Temb}(\tilde{\Sigma}^\flat(g'))$ itself has canonical singularities. We do this by extending the perfect cone, or first Voronoi, compactification of \mathcal{A}_g , which has this property by construction.

PROPOSITION 2.4. *There exist a Γ -admissible collection Σ^\flat of fans $\Sigma^\flat(g')$, for $0 \leq g' \leq g$, and a $\tilde{\Gamma}$ -admissible collection $\tilde{\Sigma}^\flat$ of fans $\tilde{\Sigma}(g')$ such that*

- (i) $|\tilde{\Sigma}^\flat(g')| = \tilde{C}(\mathbb{X})$ and $|\Sigma^\flat(g')| = C(\mathbb{X})$;
- (ii) $\tilde{\Sigma}^\flat(g')$ is $\text{GL}(\mathbb{X}) \ltimes \mathbb{X}^n$ -admissible relative to $\Sigma^\flat(g')$ for each g' ;
- (iii) $\text{Temb}(\tilde{\Sigma}^\flat(g'))$ has canonical singularities and $\tilde{\Sigma}^\flat(g')$ is equidimensional in codimension 1 over $\Sigma^\flat(g')$;
- (iv) $\sigma \times \{0\} \in \tilde{\Sigma}^\flat(g')$ for every $\sigma \in \Sigma^\flat(g')$.

Remark. The uses of the word ‘‘admissible’’ in the preamble to Proposition 2.4 and in condition (ii) there are different. The term Γ -admissible refers to the property of a collections of fans, one for each cusp, being compatible under restriction (see [AMRT10]), whereas $\mathrm{GL}(\mathbb{X}) \ltimes \mathbb{X}^n$ -admissible is a property of compatibility at each cusp separately with the projection map $X_g^n \rightarrow \mathcal{A}_g$, defined in [FC90, Definition VI.1.3].

Proof. We take $\Sigma^\flat(g')$ to be defined by the perfect cone decomposition of $C(\mathbb{X})$. This is the same as taking the cones of $\Sigma^\flat(g')$ to be the cones on the faces of the convex hull of the rank 1 forms in the closure of C with rational radical, by [BC75]. It is known to give an admissible decomposition and a polyhedral fundamental domain for the action of $\mathrm{GL}(\mathbb{X})$; see, for example, [Nam80, Chapter 8] and the references there.

We can extend this to a decomposition of $\tilde{C}(\mathbb{X})$ by taking the convex hull of all (b, ℓ) in $\tilde{C}(\mathbb{X}) \cap \mathrm{Sym}^2(\mathbb{X}) \oplus (\mathbb{X}^\vee)^n$ with rank $b = 1$. From the description of the action of $\mathrm{GL}(\mathbb{X}) \ltimes \mathbb{X}^n$, given for example in [FC90, Theorem VI.1.1], it follows immediately that $\mathrm{GL}(\mathbb{X}) \ltimes \mathbb{X}^n$ acts on the cone $\tilde{C}(\mathbb{X})$ with a polyhedral fundamental domain.

If $q = (b; \ell_1, \dots, \ell_n) = (b; \ell_j) \in \tilde{C}(\mathbb{X})$, then $b \in C(\mathbb{X})$, so we can write $b = \sum \lambda_i r_i$, where $r_i = \xi_i^t \xi_i$ is of rank 1, $\xi_i \in \mathbb{X}$ and $\lambda_i \in \mathbb{R}_+$. Since $\ell_j|_{\mathrm{rad} b} = 0$, we may write $\ell_j = b(\mu_j, \bullet)$ with $\mu_j \in \mathrm{rad} b$ (notice that $\mathrm{rad} b \supseteq \bigcup_i \mathrm{rad} r_i$, with equality if the r_i are linearly independent). Hence

$$q = (b; \ell_j) = (b; b(\mu_j, \bullet)) = \left(b; \sum_i \lambda_i r_i(\mu_j, \bullet) \right) = \sum_i \lambda_i (r_i; r_i(\mu_j, \bullet)),$$

so the cone over the convex hull of the integral q with rank 1 quadratic part b is indeed $\tilde{C}(\mathbb{X})$, and every integral point of $\tilde{C}(\mathbb{X})$ is in the convex hull.

The first of these conditions shows that the definition of $\tilde{\Sigma}^\flat$ does give an admissible collection of fans. That is, it is $\tilde{\Gamma}^n$ -invariant, and chosen for different cusps $F_{g'}$ so as to be compatible with restriction (Siegel Φ -operator) to adjacent cusps. This therefore yields a compactification $\overline{X_g^n}^\flat$. The second condition shows that the covering spaces $\mathrm{Temb}(\tilde{\Sigma}^\flat(g'))$ have canonical singularities. Finally, $\overline{X_g^n}^\flat$ is a Namikawa compactification because there are no rays of $\tilde{\Sigma}^\flat(g')$ in the interior of $\tilde{C}(\mathbb{X})$, and the ray spanned by $q = (b; \ell_j)$ projects onto the ray spanned by b . \square

Shepherd-Barron showed in [She06] (see also the correction [AS16]) that the perfect cone compactification of \mathcal{A}_g has canonical singularities for $g \geq 5$. We are not constrained to use a specific compactification; rather, we choose a suitable one, as in [Tai82, § 5]. We choose smooth subdivisions $\tilde{\Sigma}^\sharp(g')$ of the fans $\tilde{\Sigma}^\flat(g')$ (that is, toric resolutions of $\mathrm{Temb}(\tilde{\Sigma}^\flat(g'))$) in a $\tilde{\Gamma}$ -equivariant way and denote the resulting compactification by $\overline{X_g^n}^\sharp$. This is of course no longer a Namikawa compactification, nor is it smooth in general since $\tilde{\Gamma}$ is not neat. However, we have the following easy consequence of Lemma 2.3.

COROLLARY 2.5. *Suppose that $\overline{X_g^n}^\sharp$ has canonical singularities and that the action of $P(g') \cap \tilde{\Gamma}$ on $\mathrm{Temb}(\tilde{\Sigma}^\sharp(g'))$ has no quasireflections. Then $\overline{X_g^n}^\sharp$ has canonical singularities.*

Proof. It is enough to apply Lemma 2.3 to the resolutions $\mathrm{Temb}(\tilde{\Sigma}^\sharp(g')) \rightarrow \mathrm{Temb}(\tilde{\Sigma}^\flat(g'))$ for each g' . \square

3. Quotient singularities

In this section we shall verify the conditions of Corollary 2.5 for the values of g and n that concern us.

Recall that any toroidal compactification $\overline{\mathcal{A}}_g$ of \mathcal{A}_g comes with a surjective map to the Satake compactification \mathcal{A}_g^* , and that the Satake compactification has a stratification

$$\mathcal{A}_g^* = \mathcal{A}_g \sqcup \mathcal{A}_{g-1} \sqcup \cdots \sqcup \mathcal{A}_1 \sqcup \mathcal{A}_0,$$

where \mathcal{A}_0 is a point. Hence any compactification of X_g^n that dominates a Namikawa compactification has a map to \mathcal{A}_g^* , and in particular, there is a natural map $\pi: \overline{X_g^n}^\sharp \rightarrow \mathcal{A}_g^*$.

If $p \in \overline{X_g^n}^\sharp$ is such that $\pi(p) \in \mathcal{A}_{g'} \subset \mathcal{A}_g^*$, then near p the toroidal compactification is a quotient of an invariant open subset

$$\mathcal{D}^\sharp \subset \mathbb{H}_{g'} \times \mathbb{C}^{g'g''} \times \mathbb{C}^{g'n} \times \text{Temb}(\tilde{\Sigma}^\sharp) \quad (3.1)$$

(where $g' + g'' = g$) by an action of $\tilde{P}(g') \cap \tilde{\Gamma}^n$ that preserves the product structure, extending the decomposition (2.3). To determine the singularity at $p \in \overline{X_g^n}^\sharp$, we therefore have to examine the action of the stabiliser G of a preimage $\tilde{p} \in \mathcal{D}^\sharp$ of p on the tangent space $T_{\mathcal{D}^\sharp, \tilde{p}}$.

We recall some basic facts from [Tai82] and [Rei87]. Suppose that $\rho: G \rightarrow \text{GL}(m, \mathbb{C})$ is a finite-dimensional representation of a finite group G , and suppose that $h \in G$ and that $\rho(h)$ has order $k > 1$. If the eigenvalues of $\rho(h)$ are $\zeta^{a_1}, \dots, \zeta^{a_m}$ (where $\zeta = e^{2\pi i/k}$ is a primitive k th root of unity and $0 \leq a_i < k$), then the *Reid–Tai sum* of h , also called the *age* of h , is

$$\text{RT}(h) = \sum_{i=1}^m \frac{a_i}{k}.$$

The RST (Reid–Shepherd–Barron–Tai) criterion states that if $\rho(G)$ has no quasireflections, then the quotient $\mathbb{C}^m/\rho(G)$ has canonical singularities if and only if $\text{RT}(h) \geq 1$ for every $h \in G$. If $\rho(h)$ is a quasireflection, then exactly one of the a_i is non-zero, so $\text{RT}(h) < 1$. It follows that in any case if $\text{RT}(h) \geq 1$ for every $h \in G$, then the quotient has canonical singularities.

We apply this to the action of $\tilde{\gamma} \in \tilde{P}(g')$. We use the block decomposition given in (2.1) and (2.2), and the notation for submatrices in the rest of this section is taken from there.

To check that the singularity at p is canonical, it is enough to verify that G contains no quasireflection on the tangent space (which we need to do anyway in order to apply Lemma 2.3) and that $\text{RT}(\tilde{\gamma}) \geq 1$ for any non-trivial $\tilde{\gamma} \in G$.

Note that the decomposition (3.1) is G -invariant, so $\text{RT}(\tilde{\gamma})$ is the sum of the age of $\tilde{\gamma}$ restricted to each factor.

PROPOSITION 3.1. *If $g'' = 0$, then $\overline{X_g^n}$ has a canonical singularity at p and the stabiliser G of \tilde{p} has no quasireflections unless $g = 2$ and $n \leq 2$, or $g = 3$ and $n = 1$.*

Proof. Recall that we are assuming $g \geq 2$ anyway. If $g'' = 0$, then the local cover in (3.1) becomes $\mathcal{D}^\sharp = \mathbb{H}_g \times \mathbb{C}^{gn}$, which is just the covering space of X_g^n . But X_g^n has canonical singularities by [Ma21, Proposition 10.3], and there are no quasireflections by [Ma21, Lemma 7.1] for these values of g and n . \square

In view of Proposition 3.1, we may assume $g'' > 0$ for the rest of this section. In order to use the results of [Tai82], we need to verify the condition that $\tilde{\gamma}$ should act trivially on each cone of the fan (see [Tai82, §5, p. 438]).

LEMMA 3.2. *If $\tilde{\gamma}$ fixes a cusp, then it acts trivially on each cone of $\tilde{\Sigma}^\sharp(g')$.*

Proof. The eigenvectors in $\text{Sym}^2(V)$ are the rank 1 forms $f_i f_j$, where the $f_i \in \mathbb{X}^\vee \otimes \mathbb{R}$ are the real eigenvectors of u (if any), and thus there are no eigenvectors in the interior of the cone \tilde{C} . Now the result follows: if $\tilde{\gamma}$ acts non-trivially on $\sigma \in \tilde{\Sigma}^\sharp(g')$, then it acts on some subset $\{\rho_1, \dots, \rho_k\}$ of the rays spanning σ by a free permutation, and then by the Brouwer fixed-point theorem, it has an eigenvector in the closed cone spanned by $\{\rho_1, \dots, \rho_k\}$; but this eigenvector is not a generator of any of the ρ_i , so it is in the interior of \tilde{C} . \square

PROPOSITION 3.3. *Suppose that $p \in \overline{X_g^n}$ and $\pi(p) \in \mathcal{A}_{g'}$, and that $\tilde{\gamma} \neq \mathbf{1}_{n+2g}$ belongs to the stabiliser G of $\tilde{p} \in \tilde{\mathcal{D}}^\sharp$ in $\tilde{P}(g') \cap \tilde{\Gamma}^n$. Suppose that $u \neq \pm \mathbf{1}_{g''}$, or that $u = -\mathbf{1}_{g''}$ and $(g'', n) \neq (1, 1)$. Then the action of $\tilde{\gamma}$ on $T_{\mathcal{D}^\sharp, \tilde{p}}$ has $\text{RT}(\tilde{\gamma}) \geq 1$, and in particular, it is not a quasireflection.*

Proof. For this, it is enough to look at the $\text{Temb}(\tilde{\Sigma}^\sharp)$ factor. This is a toric variety with torus \mathbb{T} whose lattice of 1-parameter subgroups (the dual of the character lattice) is $\text{Sym}^2(\mathbb{X}^\vee) \times (\mathbb{X}^\vee)^n$.

If the eigenvalues of $\tilde{\gamma}$ on $V = \mathbb{X}^\vee \otimes \mathbb{C}$, which are the eigenvalues of u , are $\mu_1, \dots, \mu_{g''}$, then the eigenvalues on $\text{Sym}^2(V) \times V^n$ are $\mu_i \mu_j$ and n copies of each μ_i . The eigenvalues of $\tilde{\gamma}$ on the toric boundary component containing \tilde{p} include n copies of each μ_i and, as in [Tai82, Lemma 5.2], all the $\mu_i \mu_j$ that are different from 1. Since $u \in \text{GL}(g'', \mathbb{Z})$ is of finite order, its eigenvalues include all the primitive d th roots of unity for some degree d , and that gives $\text{RT}(\tilde{\gamma}|_V) \geq 1$ unless $d = 1$ or $d = 2$.

If $d = 1$, then $u = \mathbf{1}_{g''}$. If $d = 2$, then $\mu_i = \pm 1$ and we may assume $\mu_1 = -1$. Then if $g'' > 1$, we either have $\mu_2 = -1$, so again $\text{RT}(\tilde{\gamma}|_V) \geq 1$, or $\mu_2 = 1$, and then the eigenvalue $\mu_1 \mu_2 = -1$ occurs on $\text{Sym}^2(V)$. If $g'' = 1$, then the eigenvalue $\mu_1 = -1$ occurs n times on V^n , so $\text{RT}(\tilde{\gamma}|_{V^n}) \geq 1$ unless $g'' = n = 1$. \square

Next, we examine the action of $\tilde{\gamma}$ on the $\mathbb{C}^{ng'}$ factor. Because of Proposition 3.3, we may assume $u = \epsilon \mathbf{1}_{g''}$ with $\epsilon = \pm 1$.

For any $r \leq g$, we let $M_{r \times g}(\mathbb{C})^*$ be the set of matrices of rank r in $M_{r \times g}(\mathbb{C})$. Then the Grassmannian $\text{Gr}(r, g)$, of r -dimensional linear subspaces in \mathbb{C}^g , is $M_{r \times g}(\mathbb{C})^* / \text{GL}(r, \mathbb{C})$, with $\text{GL}(r, \mathbb{C})$ acting by right multiplication. Since \mathbb{C}^{ng} is identified with $M_{n \times g}(\mathbb{C})$ by the choice of basis e_1, \dots, e_{2g} , we may regard $\mathbb{H}_g \times \mathbb{C}^{ng}$ as a subset of $\text{Gr}(g, n+2g)$ by sending an element $(\tau, Z) \in \mathbb{H}_g \times \mathbb{C}^{ng}$ to the equivalence class of block matrices

$$\begin{bmatrix} Z \\ \tau \\ \mathbf{1}_g \end{bmatrix} \in M_{(n+2g) \times g}(\mathbb{C}) / \text{GL}(g, \mathbb{C}).$$

Recall that, in this representation, a boundary component of \mathbb{H}_g is a subset of the closure $\overline{\mathbb{H}}_g$ of \mathbb{H}_g in $\text{Gr}(g, 2g)$, so this description extends to the boundary. So the image of \tilde{p} in $\overline{\mathbb{H}}_g \times \mathbb{C}^{ng}$ may also be written in this way, with $\tau \in F_{g'} \subset \overline{\mathbb{H}}_g$ given by $\tau = \begin{pmatrix} \tau' & \omega \\ t_\omega & \tau'' \end{pmatrix}$, where $\tau' \in \mathbb{H}_{g'}$, $\omega \in M_{g' \times g''}$ and $\tau'' \in \text{Sym}^2(\mathbb{X}^\vee \otimes \mathbb{C}) = M_{g'' \times g''}^{\text{sym}}(\mathbb{C})$.

Then the action of $\tilde{\gamma}$ is given (notation from (2.1) and (2.2)) by

$$\tilde{\gamma} \cdot \begin{bmatrix} Z \\ \tau \\ \mathbf{1}_g \end{bmatrix} = \begin{bmatrix} Z + a\tau + b \\ \begin{pmatrix} A\tau' + B & Aw + m' \\ m\tau' + \epsilon^t w + n & mw + \epsilon\tau'' + M \end{pmatrix} \\ \begin{pmatrix} C\tau' + D & Cw + n' \\ 0 & \mathbf{1}_{g''} \end{pmatrix} \end{bmatrix}.$$

Because τ is preserved by γ , this simplifies to

$$\tilde{\gamma} \cdot \begin{bmatrix} Z \\ \tau \\ \mathbf{1}_g \end{bmatrix} = \begin{bmatrix} (Z + a\tau + b) \cdot N \\ \tau \\ \mathbf{1}_g \end{bmatrix},$$

where

$$N = \begin{pmatrix} (C\tau' + D)^{-1} & -(Cw + n')(C\tau' + D)^{-1} \\ 0 & \mathbf{1}_{g''} \end{pmatrix},$$

and therefore the action of $\tilde{\gamma}$ on the tangent space to \mathbb{C}^{ng} at Z is by right multiplication by N .

LEMMA 3.4. *The eigenvalues of $\tilde{\gamma}$ on \mathbb{C}^{ng} are exactly the eigenvalues of N , which are 1 and the eigenvalues of $(C\tau' + D)^{-1}$. Moreover, γ' fixes τ' ; that is, $(A\tau' + B)(C\tau' + D)^{-1} = \tau'$.*

Proof. This is immediate from the discussion above. \square

The next lemma and its corollaries apply to the usual action of $\mathrm{Sp}(2g, \mathbb{Z})$ on \mathbb{H}_g , for arbitrary g . If $\gamma = \begin{pmatrix} A_\gamma & B_\gamma \\ C_\gamma & D_\gamma \end{pmatrix} \in \mathrm{Sp}(2g, \mathbb{Z})$, then denote the eigenvalues of γ by $\Lambda = (\lambda_1, \dots, \lambda_g)$ and $\bar{\Lambda} = (\bar{\lambda}_1, \dots, \bar{\lambda}_g)$; we will regard such a sequence as a diagonal matrix.

LEMMA 3.5. *If γ fixes $\tau \in \mathbb{H}_g$, then γ is diagonalisable. Moreover, $C_\gamma\tau + D_\gamma$ is diagonalisable and has $\bar{\Lambda}$ as sequence of eigenvalues.*

Proof. For every $\beta \in \mathrm{Sp}(2g, \mathbb{R})$ and $\tau \in \mathbb{H}_g$, we set $J(\beta, \tau) = C_\beta\tau + D_\beta$. Then J is a cocycle; that is, $J(\beta_1\beta_2, \tau) = J(\beta_1, \beta_2\tau)J(\beta_2, \tau)$ for every $\beta_1, \beta_2 \in \mathrm{Sp}(2g, \mathbb{R})$ and $\tau \in \mathbb{H}_g$.

It is well known, cf. [Tai82, Lemma 4.1], that there exists an $\alpha \in \mathrm{Sp}(2g, \mathbb{R})$ such that

$$\alpha\tau = i\mathbf{1}_g \quad \text{and} \quad \alpha\gamma\alpha^{-1} = \begin{pmatrix} \delta_1 & \delta_2 \\ -\delta_2 & \delta_1 \end{pmatrix}$$

with δ_1, δ_2 real diagonal matrices and $\delta_1 + i\delta_2 \in \mathrm{U}(g, \mathbb{C})$. Obviously, γ is diagonalisable, with eigenvalues $\Lambda = \delta_1 + i\delta_2$ and $\bar{\Lambda} = \delta_1 - i\delta_2$. Now using the cocycle property, we have

$$\bar{\Lambda} = J(\alpha\gamma\alpha^{-1}, i\mathbf{1}_g) = J(\alpha, \gamma\alpha^{-1}, i\mathbf{1}_g)J(\gamma\alpha^{-1}, i\mathbf{1}_g) = J(\alpha, \gamma \cdot \tau)J(\gamma, \tau)J(\alpha^{-1}, i\mathbf{1}_g).$$

Now, since γ fixes τ and $J(\alpha^{-1}, i\mathbf{1}_g) = J(\alpha, \tau)^{-1}$, we get that $\bar{\Lambda}$ and $J(\gamma, \tau)$ are conjugate. \square

We have the following immediate corollary.

COROLLARY 3.6. *If $\gamma \in \mathrm{Sp}(2g, \mathbb{Z})$ is non-trivial and fixes $\tau \in \mathbb{H}_g$, then $C_\gamma\tau + D_\gamma$ has an eigenvalue that is not 1.*

Now we return to the singularities of \overline{X}_g^n .

PROPOSITION 3.7. *If $g'' \neq 0$, then \overline{X}_g^n has a canonical singularity at p as long as $g + n \geq 6$.*

Proof. As before, we take $\tilde{\gamma} \in \tilde{P}(g')$, fixing a point \tilde{p} , and write it using the block decomposition given in (2.1) and (2.2). Again, because of Proposition 3.3, we may assume that $u = \epsilon\mathbf{1}_{g''}$, where $\epsilon = \pm 1$ and $\epsilon = 1$ unless $g'' = n = 1$.

For any g , if $\gamma' = \mathbf{1}_{2g'}$ and $u = \mathbf{1}_{g''}$, then $\tilde{\gamma} \in \tilde{U}(g')$ (see [Nam79, Example 2.8]) and $\tilde{\gamma}$ acts trivially at the boundary $F_{g'}$. In particular, this holds if $g' = 0$ unless $g'' = n = 1$, but then $g = 1$, which is excluded.

If $\gamma' = -\mathbf{1}_{2g'}$ and $\epsilon = 1$, or $\gamma' = \mathbf{1}_{2g'}$ and $\epsilon = -1$, then there are $g - 1$ eigenvalues $\lambda_i\epsilon = -1$ on the $\mathbb{C}^{g'g''}$ factor, giving $\mathrm{RT}(\tilde{\gamma}) > 1$. If $\gamma' = -\mathbf{1}_{2g'}$ and $\epsilon = -1$, then the eigenvalues on the

$\text{Temb}(\widetilde{\Sigma}^\sharp)$ factor include n copies of ϵ , and, by Corollary 3.6, there are also n copies of -1 occurring on the \mathbb{C}^{ng} factor. Even for $n = 1$, this gives $\text{RT}(\gamma) \geq 1$. Therefore, we may assume that $\gamma' \neq \pm \mathbf{1}_{2g'}$, and thus that γ' does not act trivially on $\mathbb{H}_{g'}$.

If $g' \geq 5$ and $\gamma' \neq \pm \mathbf{1}_{2g'}$, then the contribution to $\text{RT}(\gamma)$ from γ' acting on the $\mathbb{H}_{g'}$ factor (the $F_{g'}$ factor) is already at least 1, and $\mathcal{A}_{g'}$ itself has canonical singularities: this is [Tai82, Lemma 4.5].

If $g' < 5$, then the eigenvalues not coming from the action of γ' on $\mathbb{H}_{g'}$ include g'' copies of $\epsilon\lambda_i$ on the $\mathbb{C}^{g'g''}$ factor and, by Corollary 3.6, a further n copies of $\epsilon\lambda_i^{\pm 1}$ on the \mathbb{C}^{ng} factor. If the order of $\widetilde{\gamma}$ on the tangent space at \widetilde{p} is d , then some λ_i is a non-trivial d th root of unity, and so this gives a contribution of at least $(n + g'')/d$ to $\text{RT}(\widetilde{\gamma})$.

Moreover, according to [Tai82, Lemma 4.4] we may assume $d \leq 6$, and in each case the action of γ' contributes at least g'/d to $\text{RT}(\widetilde{\gamma})$, so in any case we have $\text{RT}(\widetilde{\gamma}) \geq (g' + n + g'')/d \geq (n + g)/6$. So if $n + g \geq 6$, we are done. \square

The condition $n + g \geq 6$ cannot be strengthened: it is needed if $g' = 1$ and $d = 6$.

Theorem 2.2 now follows immediately.

4. Slope of \mathcal{A}_g

We assume $g > 1$ throughout since the case $g = 1$ can be reduced to the case of $\mathcal{M}_{1,n+1}$, which is solved in [Bel98] and [BF06].

We shall construct differential forms by using Siegel modular forms, so we begin with some elementary definitions concerning them.

DEFINITION 4.1. A *modular form of weight k* is a holomorphic function $f: \mathbb{H}_g \rightarrow \mathbb{C}$ on the Siegel upper half-plane $\mathbb{H}_g = \{Z \in M_{g \times g}(\mathbb{C}) \mid Z = {}^t Z, \text{Im } Z > 0\}$ such that

$$f(\gamma \cdot \tau) = \det(C\tau + D)^k f(\tau) \quad \text{for any } \gamma \in \text{Sp}(2g, \mathbb{Z}).$$

Note that we need no extra condition at infinity when $g > 1$.

A Siegel modular form has a Fourier expansion

$$f(\tau) = \sum_T a(T) \exp(\pi i \text{tr}(T\tau)),$$

where the sum runs over all even integral symmetric matrices T .

DEFINITION 4.2. If f is a Siegel modular form, the *vanishing of f at the boundary* is

$$b := \frac{1}{2} \min \{x^t T x \mid a(T) \neq 0, x \in \mathbb{Z}^g \setminus \{0\}\}.$$

If $b > 0$, that is, if $a(T) \neq 0$ implies $T > 0$, we say that f is a *cusp form*.

We recall, mainly from [Mum83a], some facts about $\mathcal{A}_g = \text{Sp}(2g, \mathbb{Z}) \backslash \mathbb{H}_g$ and its compactifications. Modular forms of weight 1 determine a \mathbb{Q} -line bundle L , the Hodge line bundle. The Satake compactification \mathcal{A}_g^* is Proj of the ring of modular forms, and the Mumford partial compactification \mathcal{A}'_g is the blow-up of $\mathcal{A}_g \sqcup \mathcal{A}_{g-1} \subset \mathcal{A}_g^*$ along \mathcal{A}_{g-1} . Every toroidal compactification of \mathcal{A}_g dominates \mathcal{A}_g^* and contains \mathcal{A}'_g as a Zariski open subset: the toroidal compactifications differ from one another only above the deeper strata $\mathcal{A}_{g'}$ for $g' < g - 1$.

If $g \geq 2$, then $\text{Pic}(\mathcal{A}'_g) \otimes \mathbb{Q} = \mathbb{Q}\lambda \oplus \mathbb{Q}\delta$, where λ is the class of L and δ is the class of the boundary divisor $\Delta_{\mathcal{A}'}$, the proper transform of $\mathcal{A}_{g-1} \subset \mathcal{A}_g^*$. This is proved in [Mum83a,

Corollary 1.6] for $g \geq 4$, and for $g = 2$ and $g = 3$ it follows from the studies of the Chow ring in [Mum83b] and [Fab90], respectively; see [Hul00].

DEFINITION 4.3. The *slope* of an effective divisor $E = a\lambda - b\delta$ on \mathcal{A}'_g with $a, b > 0$ is defined to be $s(E) = a/b$. In particular, if E is the zero divisor of a cusp form f of weight k and vanishing order b , then $s(E) = k/b$.

The Kodaira dimension of the Kuga varieties X_g^n is related to the transcendence degree of the field generated by cusp forms of slope less than or equal to $g + n + 1$; see for example [Ma21, Theorem 1.3].

DEFINITION 4.4. The *minimal slope* $s_{\min}(g)$ is the infimum of the slopes of all effective divisors on \mathcal{A}'_g .

An upper bound $s_{\min}(g)$ is provided in small genera by the Andreotti–Mayer divisor N_0 , the locus of principally polarised abelian varieties with singular theta divisor [AM67]. The divisor N_0 has two components: Θ_{null} , the locus where the theta divisor has a singular point of order 2, and N'_0 , the locus where the theta divisor has a singular point not of order 2. The classes of Θ_{null} , for $g \geq 1$, and N'_0 , for $g \geq 4$, are computed in [Mum83a]:

$$[\Theta_{\text{null}}] = 2^{g-2}(2^g + 1)\lambda - 2^{2g-5}\delta,$$

$$[N'_0] = \left(\frac{(g+1)!}{4} + \frac{g!}{2} - 2^{g-3}(2^g + 1) \right) \lambda - \left(\frac{(g+1)!}{24} - 2^{2g-6} \right) \delta.$$

For $g \leq 3$ the minimal slope is achieved at $[\Theta_{\text{null}}]$, giving the values $s_{\min}(1) = 12$, $s_{\min}(2) = 10$ and $s_{\min}(3) = 9$. For $g = 4$ we have $s_{\min}(4) = 8$, achieved by $s([N'_0])$, and for $g = 5$, we have $s_{\min}(5) = 54/7$, also achieved by $s([N'_0])$; see [Sal92] and [FGSV14], respectively. For $2 \leq g \leq 4$ the divisors that minimise the slope are rigid. For $g = 6$ we have $s_{\min}(6) \leq 7$; see [DSS21]. However, the fact that $s_{\min}(6) \leq s([N'_0]) = 550/73 < 8$ will suffice for our purposes. For $g \geq 7$ we have $s_{\min}(g) < g + 1$ by [Tai82] and [Mum83a].

THEOREM 4.5. Suppose $g \geq 2$ and that $\overline{X_g^n}$ is a Namikawa compactification of X_g^n with canonical singularities. Then

- (i) $\kappa(X_g^n) = \frac{1}{2}g(g+1)$ if $s_{\min}(g) < g + n + 1$;
- (ii) $\kappa(X_g^n) = 0$ if $s_{\min}(g) = s(D) = g + n + 1$ and D is rigid;
- (iii) $\kappa(X_g^n) = -\infty$ if $s_{\min}(g) > g + n + 1$ (even if the singularities are not canonical).

Proof. The first case (what one might call relatively general type) follows easily from [Ma21, Proposition 9.2]. Pulling back along $f: \overline{X_g^n} \xrightarrow{\#} \overline{\mathcal{A}}_g$, this implies $kK_X \geq f^*(k(g+n+1)L - k\Delta_{\mathcal{A}})$ for sufficiently divisible k . So it is enough to show that the \mathbb{Q} -divisor $(g+n+1)L - \Delta_{\mathcal{A}}$ is big; however, since it has slope strictly greater than $s_{\min}(g)$, it is in the interior of the effective cone and can therefore be written as the sum of an effective divisor and an ample divisor.

In the second case, the same argument shows that K_X is effective since it dominates the pullback of the effective divisor $(g+n+1)L - \Delta_{\mathcal{A}}$. Therefore, $\kappa(X_g^n) \geq 0$. On the other hand, if some multiple of K_X moves, then so does some multiple of $(g+n+1)L - \Delta_{\mathcal{A}}$, which is to say that some multiple of D moves, but D is rigid.

In the third case, if $K_X \geq 0$, then $f_*(K_X) \geq 0$, but $s(f_*(K_X)) = g + n + 1 < s_{\min}(g)$. So K_X is not effective, and $\kappa(X_g^n) = -\infty$. \square

Theorem 1 follows immediately from this.

ACKNOWLEDGEMENTS

The authors would like to thank G. Farkas, S. Ma and N. I. Shepherd-Barron for useful discussions on this topic, and the referee for asking about the relation with $\mathcal{M}_{g,n}$.

REFERENCES

AM67 A. Andreotti and A. L. Mayer, *On period relations for abelian integrals on algebraic curves*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (3) **21** (1967), no. 2, 189–238.

AS16 J. Armstrong and N. I. Shepherd-Barron, *The singularities of \mathcal{A}_g^P* , 2016, [arXiv:1604.05943](https://arxiv.org/abs/1604.05943).

AMRT10 A. Ash, D. Mumford, M. Rapoport and Y.-S. Tai, *Smooth compactification of locally symmetric varieties*, 2nd ed., Cambridge Math. Lib. (Cambridge Univ. Press, Cambridge, 2010); [doi:10.1017/CBO9780511674693](https://doi.org/10.1017/CBO9780511674693).

BC75 E. S. Barnes and M. J. Cohn, *On the inner product of positive quadratic forms*, J. London Math. Soc. (2) **12** (1975), no. 1, 32–36; [doi:10.1112/jlms/s2-12.1.32](https://doi.org/10.1112/jlms/s2-12.1.32).

Bel98 P. Belorousski, *Chow rings of moduli spaces of pointed elliptic curves*, Ph.D. thesis, 1998, University of Chicago, available at <https://www.proquest.com/docview/304462671>.

BF06 G. Bini and C. Fontanari, *Moduli of curves and spin structures via algebraic geometry*, Trans. Am. Math. Soc. **358** (2006), no. 7, 3207–3217; [doi:10.1090/S0002-9947-06-03838-4](https://doi.org/10.1090/S0002-9947-06-03838-4).

BFV12 G. Bini, C. Fontanari and F. Viviani, *On the birational geometry of the universal Picard variety*, Int. Math. Res. Not. IMRN (2012), no. 4, 740–780; [doi:10.1093/imrn/rnr045](https://doi.org/10.1093/imrn/rnr045).

Cap94 L. Caporaso, *A compactification of the universal Picard variety over the moduli space of stable curves*, J. Amer. Math. Soc. **7** (1994), no. 3, 589–660; [doi:10.2307/2152786](https://doi.org/10.2307/2152786).

CKV17 S. Casalaina-Martin, J. L. Kass and F. Viviani, *The singularities and birational geometry of the compactified universal Jacobian*, Algebr. Geom. **4** (2017), no. 3, 353–393; [doi:10.14231/AG-2017-020](https://doi.org/10.14231/AG-2017-020).

DSS21 M. Dittmann, R. Salvati Manni and N. R. Scheithauer, *Harmonic theta series and the Kodaira dimension of \mathcal{A}_6* , Algebra Number Theory **15** (2021), no. 1, 271–285; [doi:10.2140/ant.2021.15.271](https://doi.org/10.2140/ant.2021.15.271).

Fab90 C. Faber, *Chow rings of moduli spaces of curves. I. The Chow ring of $\overline{\mathcal{M}}_3$* , Ann. of Math. (2) **132** (1990), no. 2, 331–419; [doi:10.2307/1971525](https://doi.org/10.2307/1971525).

FC90 G. Faltings and C.-L. Chai, *Degeneration of abelian varieties*, Ergeb. Math. Grenzgeb. (3), vol. 22 (Springer-Verlag, Berlin, 1990); [doi:10.1007/978-3-662-02632-8](https://doi.org/10.1007/978-3-662-02632-8).

FGSV14 G. Farkas, S. Grushevsky, R. Salvati Manni and A. Verra, *Singularities of theta divisors and the geometry of \mathcal{A}_5* , J. Eur. Math. Soc. (JEMS) **16** (2014), no. 9, 1817–1848; [doi:10.4171/JEMS/476](https://doi.org/10.4171/JEMS/476).

FV13 G. Farkas and A. Verra, *The classification of universal Jacobians over the moduli space of curves*, Comment. Math. Helv. **88** (2013), no. 3, 587–611; [doi:10.4171/CMH/297](https://doi.org/10.4171/CMH/297).

FV16 ———, *The universal abelian variety over \mathcal{A}_5* , Ann. Sci. Éc. Norm. Supér. (4) **49** (2016), no. 3, 521–542; [doi:10.24033/asens.2289](https://doi.org/10.24033/asens.2289).

Hul00 K. Hulek, *Nef divisors on moduli spaces of abelian varieties*, in *Complex analysis and algebraic geometry* (de Gruyter, Berlin, 2000), 255–274; [doi:10.1515/9783110806090-012](https://doi.org/10.1515/9783110806090-012).

Kaw85 Y. Kawamata, *Minimal models and the Kodaira dimension of algebraic fiber spaces*, J. reine angew. Math. **363** (1985), 1–46; [doi:10.1515/crll.1985.363.1](https://doi.org/10.1515/crll.1985.363.1).

KT21 H. Keneshlou and F. Tanturri, *On the unirationality of moduli spaces of pointed curves*, Math. Z. **299** (2021), no. 3–4, 1973–1986; [doi:10.1007/s00209-021-02741-7](https://doi.org/10.1007/s00209-021-02741-7).

Kug18 M. Kuga, *Kuga Varieties. Fiber varieties over a symmetric space whose fibers are abelian varieties.*, CTM. Class. Top. Math., vol. 9 (Higher Education Press, Beijing, 2018).

Ma21 S. Ma, *Universal abelian variety and Siegel modular forms*, Algebra Number Theory **15** (2021), no. 8, 2089–2122; [doi:10.2140/ant.2021.15.2089](https://doi.org/10.2140/ant.2021.15.2089).

Mum83a D. Mumford, *On the Kodaira dimension of the Siegel modular variety*, Algebraic geometry—open problems (Ravello, 1982), Lecture Notes in Math., vol. 997 (Springer, Berlin, 1983), 348–375; [doi:10.1007/BFb0061652](https://doi.org/10.1007/BFb0061652).

Mum83b _____, *Towards an enumerative geometry of the moduli space of curves*, in *Arithmetic and geometry, Vol. II*, Progr. Math., vol. 36 (Birkhäuser, Boston, MA, 1983), 271–328; [doi:10.1007/978-1-4757-9286-7_12](https://doi.org/10.1007/978-1-4757-9286-7_12).

Nam79 Y. Namikawa, *Toroidal degeneration of abelian varieties. II*, Math. Ann. **245** (1979), no. 2, 117–150; [doi:10.1007/BF01428801](https://doi.org/10.1007/BF01428801).

Nam80 _____, *Toroidal compactification of Siegel spaces*, Lecture Notes in Math., vol. 812 (Springer, Berlin, 1980); [doi:10.1007/BFb0091051](https://doi.org/10.1007/BFb0091051).

Rei87 M. Reid, *Young person’s guide to canonical singularities*, Algebraic geometry, Part 1, Bowdoin, 1985 (Brunswick, Maine, 1985), Proc. Symp. Pure Math., vol. 46 (Amer. Math. Soc., Providence, RI, 1987), 345–414; [doi:10.1090/pspum/046.1/927963](https://doi.org/10.1090/pspum/046.1/927963).

Sal92 R. Salvati Manni, *Modular forms of the fourth degree. Remark on the paper: “Slopes of effective divisors on the moduli space of stable curves” by J. D. Harris and I. Morrison*, Classification of irregular varieties (Trento, 1990), Lecture Notes in Math., vol. 1515 (Springer, Berlin, 1992), 106–111; [doi:10.1007/BFb0098340](https://doi.org/10.1007/BFb0098340).

She06 N. I. Shepherd-Barron, *Perfect forms and the moduli space of abelian varieties*, Invent. Math. **163** (2006), no. 1, 25–45; [doi:10.1007/s00222-005-0453-0](https://doi.org/10.1007/s00222-005-0453-0).

Tai82 Y.-S. Tai, *On the Kodaira dimension of the moduli space of abelian varieties*, Invent. Math. **68** (1982), no. 3, 425–439; [doi:10.1007/BF01389411](https://doi.org/10.1007/BF01389411).

Ver08 A. Verra, *On the universal principally polarized abelian variety of dimension 4*, Curves and abelian varieties (Athens GA, 2007), Contemp. Math., vol. 465 (Amer. Math. Soc., Providence, RI, 2008), 253–274; [doi:10.1090/conm/465/09106](https://doi.org/10.1090/conm/465/09106).

Flora Poon wkpoon@ncts.ntu.edu.tw

Mathematics Division, National Center for Theoretical Sciences, No. 1, Sec. 4, Roosevelt Rd., Taipei City, Taiwan

Riccardo Salvati Manni salvati@mat.uniroma1.it

Dipartimento di Matematica “Guido Castelnuovo”, Università di Roma “La Sapienza”, P.le Aldo Moro 5, 00185 Roma, Italy

Gregory Sankaran G.K.Sankaran@bath.ac.uk

Department of Mathematical Sciences, University of Bath, Bath BA2 7AY, UK